

**Problem 1.21**

Show that a number  $x \in [0, 1]$  has more than one  $p$ -adic expansion if and only if  $x = \sum_{k=1}^n a_k/p^k$  for some  $n$  where  $a_n \neq 0$ . Show also that in this case  $x$  has exactly one other expansion,

$$x = \sum_{k=1}^{n-1} a_k/p^k + \frac{a_n - 1}{p^n} + \sum_{k=n+1}^{\infty} \frac{p-1}{p^k}. \quad (1)$$

Also, characterize the numbers in  $[0, 1]$  with repeating and eventually repeating  $p$ -adic expansions.

**Solution**

Suppose  $x$  has two different expansions,

$$\begin{aligned} x &= \sum_{k=1}^{\infty} \frac{a_k}{p^k} \\ &= \sum_{k=1}^{\infty} \frac{b_k}{p^k}. \end{aligned}$$

Let  $N$  be the index in which they first differ, and without loss of generality assume that  $a_N > b_N$ . We will show that  $a_N = b_N + 1$ ,  $a_n = 0$  for  $n > N$ , and  $b_n = p - 1$  for  $n > N$ . This will prove that if  $x$  has two different expansions, then one must be a terminating expansion, and that the only other expansion is the one of the form (1).

Let  $y = \sum_{k=1}^{N-1} \frac{b_k}{p^k}$ . Then

$$\begin{aligned} x &\leq \sum_{k=1}^{N-1} \frac{b_k}{p^k} + \frac{b_N}{p^N} + \sum_{k=N+1}^{\infty} \frac{p-1}{p^k} \\ &= y + \frac{b_N}{p^N} + \frac{p-1}{p^{N+1}} \frac{p}{p-1} \\ &= y + \frac{b_N}{p^N} + \frac{1}{p^N} \end{aligned}$$

with strict inequality unless  $b_n = p - 1$  for all  $n > N$ . Similarly,

$$\begin{aligned} x &\geq \sum_{k=1}^{N-1} \frac{a_k}{p^k} + \frac{a_N}{p^N} + \sum_{k=N+1}^{\infty} \frac{0}{p^k} \\ &= y + \frac{a_N}{p^N} \end{aligned}$$

with strict inequality unless  $a_n = 0$  for  $n > N$ . These inequalities together imply

$$\frac{a_N}{p^N} \leq \frac{b_N + 1}{p^N}$$

and hence  $a_N \leq b_N + 1$  (with strict inequality unless  $b_n = p - 1$  and  $a_n = 0$  for  $n > N$ ). But  $a_N \geq b_N + 1$  since  $a_N > b_N$  and since  $a_N$  and  $b_N$  are integers. Hence  $a_N = b_N + 1$  and  $b_n = p - 1$  and  $a_n = 0$  for  $n > N$ .

If  $x$  has a terminating  $p$ -adic expansion, then  $x$  is of the form

$$x = \frac{a}{p^N}$$

where  $N \in \mathbb{N}$  and  $0 \leq a \leq p^N$ .

If  $x$  has a repeating  $p$ -adic expansion, then  $x$  is of the form

$$\frac{a}{p^N - 1}$$

where  $N \in \mathbb{N}$  and where  $0 \leq a \leq p^N - 1$ .

If  $x$  has an eventually repeating  $p$ -adic expansion, then  $x$  can be decomposed into a terminating and a repeating part. So  $x$  is of the form

$$x = \frac{a}{p^N} + \frac{1}{p^N} \frac{b}{p^M - 1}$$

where  $N, M \in \mathbb{N}$ ,  $0 \leq a < p^N$ , and  $0 \leq b \leq p^M - 1$ . This is easily seen to be the same as

$$x = \frac{d}{p^N(p^M - 1)}$$

for some  $d$  with  $0 \leq d \leq p^N(p^M - 1)$ . It is perhaps surprising that every rational admits an expression of this form.